

ASSESSMENT OF ECOLOGICAL AND AGRICULTURAL RISKS OF
HEAVY METAL POLLUTION OF SOILS IN THE VICINITY
OF KAPAN CITY, ARMENIA

G. A. GEVORGYAN ^{1*}, K. A. GHAZARYAN ², H. S. MOVSESYAN ^{2**}

¹ *Scientific Center of Zoology and Hydroecology of NAS RA*

² *Chair of Ecology and Nature Protection of YSU, Armenia*

Ecological and agricultural risks of heavy metal pollution of soils in risky areas around Kapan City (RA) were investigated. Soil samples were collected in June 2013 and analyzed for Cr, Mn, Ni, Cu, Zn, As, Mo, Cd, Pb, Co, Hg by the mass spectrometric method. The investigations showed that the soils around Kapan copper-molybdenum combine and Geghanush tailing dump were significantly polluted with heavy metals, which may have posed serious risks not only to soil biological health, but also to agricultural production.

Keywords: mining activity, heavy metals, soil, ecological and agricultural risks.

Introduction. Soil is the key part of the Earth system as it controls the hydrological, erosional, biological and geochemical cycles. The soil system also offers goods, services, and resources to Humankind [1, 2]. Soil pollution is defined as a phenomenon characterized by the loss of structural and biological properties by the soil layers as a result of numerous human and natural factors [3]. Heavy metal pollution of soil has become a critical environmental concern due to its potential adverse ecological effects. Heavy metals occur naturally at low concentrations in soils. However, they are considered as soil contaminants due to their widespread occurrence, as well as their acute and chronic toxicity. These metals are extremely persistent in the environment. They are non-biodegradable, non-thermo-degradable and thus readily accumulate to toxic levels [4]. Soil contamination by heavy metals can lead to changes in soil characteristics and limit productive and environmental functions [5]. Since heavy metals do not break down, they might affect the biosphere for a long time [4]. Polluted soils are no longer appropriate for agricultural production, because they are unable to produce healthy food [5]. In recent years, with the development of global economy, both type and content of heavy metals in soil caused by human activities have gradually increased, resulting in the deterioration of the environment [6].

Mining and smelting operations are important causes of heavy metal contamination in the environment due to activities such as mineral excavation, ore transportation, smelting, refining and the disposal of tailings and wastewater

* E-mail: gev_gor@mail.ru

** E-mail: hasmikmov@ysu.am

around mines. Mining and metallurgical industries are highly developed in the RA, particularly in Syunik Marz and Lori Marz. Kapan Region (Syunik Marz, South-Eastern part of RA), where Kapan and Zangezur copper-molybdenum combines are operating, is considered as the main metallurgical industrial centre in Armenia. Human activities in this territory are mainly expressed by heavy metal emission into the environment [7]. Therefore, for conserving the soil resources and ensuring environmental safety in this territory, the investigation of heavy metal pollution in soils and of related environmental risks is urgently required.

The aim of the article was to investigate and assess the ecological and agricultural risks of heavy metal pollution of soils in risky areas around Kapan City.

Materials and Methods. The soils in the vicinity of Kapan City were investigated in June 2013. The investigated soils belong to the type of mountain cambisol. Observation sites were selected around Kapan copper-molybdenum combine (KCMC) (№№ 1–6) and Geghanush tailing dump (GTD) (№№ 7–12). A control site was selected 4 km away from KCMC.

The soil samples were obtained from a depth of 0–20 cm and transferred into well labeled polyethylene bags for storage and laboratory analysis. The collected samples were air-dried at room temperature. The dried samples were grounded into powder by a laboratory mortar and pestle, sieved with 1 mm mesh and stored in an air tight container prior to analysis. The soil samples were digested by the Aqua Regia (conc. HCl and conc. HNO₃, 3:1) digestion method [8]. The digested soil samples were analyzed for heavy metals (Cr, Mn, Ni, Cu, Zn, As, Mo, Cd, Pb, Co, Hg) by using ELAN 9000 inductively coupled plasma mass spectrometer (ICP–MS) [9].

The ecological risks of heavy metals in the soils were assessed according to the Potential ecological risk index (PERI) method [6], calculated as follows:

$$C_r^i = C_s^i / C_n^i, \quad E_r^i = C_r^i T_r^i, \quad \text{RI} = \sum E_r^i, \quad (1)$$

where C_r^i , C_s^i , C_n^i are the pollution factor; the measured concentration and the background concentration of a single element in soil, respectively; E_r^i is the PERI of a single element; T_r^i is the toxic response factor for a single element; RI is the comprehensive PERI. The classification of RI categories according to the PERI values is presented in Tab. 1 [10].

Table 1

The adjusted grading standard of potential ecological risk of heavy metals in soil

| E_r^i | Pollution degree | RI | Risk level | Risk degree |
|------------------------|------------------|----------------------------|------------|-------------|
| $E_r^i < 30$ | slight | RI < 40 | A | slight |
| $30 \leq E_r^i < 60$ | medium | $40 \leq \text{RI} < 80$ | B | medium |
| $60 \leq E_r^i < 120$ | strong | $80 \leq \text{RI} < 160$ | C | strong |
| $120 \leq E_r^i < 240$ | very strong | $160 \leq \text{RI} < 320$ | D | very strong |
| $E_r^i \geq 240$ | extremely strong | RI ≥ 320 | – | |

Individual heavy metal pollution degree for agricultural production on soil (Pollution index (PI)) was assessed by the following equation: $\text{PI} = C_i / S_i$, where C_i is the measured concentration of heavy metal i in soil; S_i is the maximum permissible concentration of heavy metal i for agricultural production in soil [11, 12].

The PI of each metal is classified into five pollution categories: non-pollution ($PI < 1$), low level of pollution ($1 \leq PI < 2$), moderate level of pollution ($2 \leq PI < 3$), strong level of pollution ($3 \leq PI < 5$), very strong level of pollution ($PI > 5$) [11].

Integrated heavy metal pollution degree for agricultural production on soil was evaluated by the Nemerow integrated pollution index (NIPI):

$$NIPI = \sqrt{PI_{avg}^2 + PI_{max}^2} / 2, \tag{2}$$

where PI_{avg} is the average value of the single pollution indices of all heavy metals; PI_{max} is the maximum value of the single pollution indices of all heavy metals [11, 13]. NIPI is classified by the following pollution categories: non-pollution ($NIPI \leq 0.7$), warning line of pollution ($0.7 < NIPI \leq 1$), low level of pollution ($1 < NIPI \leq 2$), moderate level of pollution ($2 < NIPI \leq 3$), high level of pollution ($NIPI > 3$) [11].

Results and Discussion. The results of the study of heavy metals content in the investigated areas showed that the concentrations of different heavy metals in all the investigated observation sites exceeded the background (control) level, which indicated that heavy metal concentrations in the soils were conditioned by both natural and anthropogenic sources (Tab. 2). The main source of the heavy metal pollution of the soils in this area is KCMC activity.

Table 2

Heavy metal concentrations (mg/kg) in the soils around KCMC (№№ 1–6) and GTD (№№ 7–12)

| Sampling site | Cr | Mn | Ni | Cu | Zn | As | Mo | Cd | Pb | Co | Hg |
|---------------|-----------|--------------|------------|------------|------------|----------|---------|---------|----------|-----------|---------|
| №№ 1–6 | 17.5–99.5 | 813.2–1557.6 | 26.4–118.5 | 32.6–72.8 | 66.0–127.1 | 6.9–22.5 | 0.3–1.1 | 0.2–0.5 | 4.0–22.4 | 11.7–28.1 | 0.1–0.3 |
| №№ 7–12 | 18.6–47.4 | 854.4–1611.5 | 20.5–56.1 | 37.8–100.4 | 56.2–97.7 | 2.7–7.8 | 0.1–0.5 | 0.1–0.2 | 2.6–7.1 | 17.0–30.1 | 0.1–0.2 |
| Control | 43.2 | 527.7 | 26.4 | 9.0 | 75.5 | 7.5 | 0.2 | 0.2 | 10.3 | 10.3 | 0.1 |

Table 3

Individual and integrated metal potential ecological risk index values in the soils

| Sampling site | $E_r(\text{Cr})$ | $E_r(\text{Mn})$ | $E_r(\text{Ni})$ | $E_r(\text{Cu})$ | $E_r(\text{Zn})$ | $E_r(\text{As})$ | $E_r(\text{Mo})$ | $E_r(\text{Cd})$ | $E_r(\text{Pb})$ | $E_r(\text{Co})$ | RI |
|---------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------|
| KCMC | 0.81–4.60 | 1.54–2.95 | 5.00–22.45 | 18.15–40.56 | 0.87–1.68 | 9.17–30.07 | 1.39–6.33 | 28.75–65.00 | 1.96–10.93 | 5.10–12.30 | 86.31–171.41 |
| GTD | 0.86–2.20 | 1.62–3.05 | 3.89–10.62 | 21.09–55.98 | 0.74–1.29 | 3.66–10.39 | 0.67–2.94 | 15.00–28.75 | 1.29–3.47 | 7.45–13.17 | 60.95–108.35 |

For assessing the biological health risks of heavy metal pollution in the investigated soils, the PERI values, which represent the sensitivity of various biological communities to harmful elements and illustrate the potential ecological risk, were applied (Tab. 3) [14]. According to them, the soils were significantly polluted with heavy metals (see the pollution degree in Tab. 4). All of this may have caused soil’s biological health risks, the levels of which are illustrated in Tab. 5. The highest health risks to biological communities in the soils around KCMC may have been posed by Cd (slight-strong) and around GTD by Cu (mainly medium) (Tab. 5).

According to the values, individual metal pollution degree and related potential health risks to biological communities were in the order of $Cd \leq Cu \leq As \leq Ni \leq Co \leq Pb \leq Mo \leq Cr \leq Mn \leq Zn$ (soils around KCMC) and $Cu \leq Cd \leq Co \leq Ni \leq As \leq Mn \leq Pb \leq Mo \leq Cr \leq Zn$ (soils around GTD) (Tab. 3).

Table 4

Individual and integrated metal pollution degree and potential ecological risk levels for the biological health of the soils

| Sampling site | $E_r(\text{Cr})$ | $E_r(\text{Mn})$ | $E_r(\text{Ni})$ | $E_r(\text{Cu})$ | $E_r(\text{Zn})$ | $E_r(\text{As})$ | $E_r(\text{Mo})$ | $E_r(\text{Cd})$ | $E_r(\text{Pb})$ | $E_r(\text{Co})$ | RI | |
|---------------------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|----|---|
| Kapan copper-molybdenum combine | 1 | A | A | A | B | A | A | A | B | A | A | D |
| | 2 | A | A | A | A | A | A | A | B | A | A | C |
| | 3 | A | A | A | A | A | A | A | B | A | A | C |
| | 4 | A | A | A | A | A | A | A | B | A | A | C |
| | 5 | A | A | A | B | A | A | A | A | A | A | C |
| | 6 | A | A | A | A | A | B | A | C | A | A | C |
| Geghanush tailing dump | 7 | A | A | A | B | A | A | A | A | A | A | C |
| | 8 | A | A | A | B | A | A | A | A | A | A | C |
| | 9 | A | A | A | B | A | A | A | A | A | A | C |
| | 10 | A | A | A | B | A | A | A | A | A | A | C |
| | 11 | A | A | A | A | A | A | A | A | A | A | B |
| | 12 | A | A | A | B | A | A | A | A | A | A | C |

A – slight; B – medium; C – strong; D – very strong.

Table 5

Individual and integrated heavy metal pollution degree for agricultural production of the soils

| Sampling site | PI_{Cr} | PI_{Mn} | PI_{Ni} | PI_{Cu} | PI_{Zn} | PI_{As} | PI_{Mo} | PI_{Cd} | PI_{Pb} | PI_{Hg} | NIPI |
|---------------------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------|
| Kapan copper-molybdenum combine | 1 | | | | | | | | | | |
| | 2 | | | | | | | | | | |
| | 3 | | | | | | | | | | |
| | 4 | | | | | | | | | | |
| | 5 | | | | | | | | | | |
| | 6 | | | | | | | | | | |
| Geghanush tailing dump | 7 | | | | | | | | | | |
| | 8 | | | | | | | | | | |
| | 9 | | | | | | | | | | |
| | 10 | | | | | | | | | | |
| | 11 | | | | | | | | | | |
| | 12 | | | | | | | | | | |

Level of pollution: □ – non-pollution, ◻ – warning, ◼ – low, ◼ – moderate level.

Table 6

PI and NIPI values for agricultural production of the soils

| Sampling site | PI_{Cr} | PI_{Mn} | PI_{Ni} | PI_{Cu} | PI_{Zn} | PI_{As} | PI_{Mo} | PI_{Cd} | PI_{Pb} | PI_{Hg} | NIPI |
|---------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|-----------|
| KCMC | 0.19–1.11 | 0.54–1.04 | 0.33–1.48 | 0.25–0.55 | 0.30–0.58 | 0.69–2.25 | 0.00–0.01 | 0.12–0.26 | 0.06–0.35 | 0.10–0.15 | 0.62–1.67 |
| GTD | 0.21–0.53 | 0.57–1.07 | 0.26–0.70 | 0.29–0.76 | 0.26–0.44 | 0.27–0.78 | 0.00–0.00 | 0.08–0.12 | 0.04–0.11 | 0.06–0.10 | 0.47–0.82 |

The results of the study showed that heavy metal pollution degree in the investigated soils may have adversely affected not only soil biological health, but also agricultural production (Tabs. 4, 5). The highest risks to agricultural production of the soils around KCMC may have been posed by As and around GTD by Mn (Tab. 5). The investigated heavy metals can be ranked by individual metal pollution degree for agricultural production as follows: $\text{As} \leq \text{Ni} \leq \text{Mn} \leq \text{Cr} \leq \text{Zn} \leq \text{Cu} \leq \text{Pb} \leq \text{Cd} \leq \text{Hg} \leq \text{Mo}$

in the soils around KCMC and $Mn \leq As \leq Ni \leq Cr \leq Cu \leq Zn \leq Cd \leq Hg \leq Pb \leq Mo$ in the soils around GTD (Tab. 6). It is necessary to mention that some heavy metals pollution degree (for example, Cu and Cd levels), being the major stressor of soil's biological communities wasn't dangerous for agricultural production (Tabs. 4, 5). All of this indicates that individual heavy metal pollution degree in soil should be estimated for each of vulnerable components individually, which will enable to precisely understand all the related environmental risks.

Conclusion. In general, it is possible to state that KCMC activity caused significant heavy metal pollution in the soils. Such a heavy metal pollution degree not only may have posed serious soil's biological health risks, but also may have been dangerous for agricultural production. To mitigate such environmental risks, responsible authorities need to develop and implement a new policy of mining operation, which will take into consideration not only economic benefits, but also environmental security.

This work was supported by the SCS MES RA, in the frame of the research project SCS RA № 13-4C202.

Received 31.07.2017

REFERENCES

1. **Adamcova D.** et al. Soil Contamination in Landfills: a Case Study of a Landfill in Czech Republic. // *Solid Earth*, 2016, v. 7, № 1, p. 239–247.
2. **Adugna A.** et al. Soil Erosion Assessment and Control in Northeast Wollega, Ethiopia. // *Solid Earth Discussions*, 2015, v. 7, № 4, p. 3511–3540.
3. **Gangadhar Z.S.** Environmental Impact Assessment on Soil Pollution Issue About Human Health. // *International Research Journal of Environmental Sciences*, 2014, v. 3, № 11, p. 78–81.
4. **Karishma B., Prasad S.H.** Effect of Agrochemicals Application on Accumulation of Heavy Metals on Soil of Different Land Uses with Respect to Its Nutrient Status. // *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 2014, v. 8, № 7, p. 46–54.
5. **Angelovicova L., Fazekasova D.** Contamination of the Soil and Water Environment by Heavy Metals in the Former Mining Area of Rudnany (Slovakia). // *Soil and Water Research*, 2014, v. 9, № 1, p. 18–24.
6. **Hakanson L.** An Ecological Risk Index for Aquatic Pollution Control a Sedimentological Approach. // *Water Research*, 1980, v. 14, № 8, p. 975–1001.
7. RA Government Decision About the Approvement of the Order of the Evaluation of Economic Activities – Caused Impact on Soil Resources, 2005 (in Armenian). <http://www.arlis.am/DocumentView.aspx?DocID=13401>
8. **Sarala T.D., Sabitha M.A.** Calculating Integrated Pollution Indices for Heavy Metals in Ecological Geochemistry Assessment Near Sugar Mill. // *J. of Research in Biology*, 2012, v. 2, № 5, p. 489–498.
9. **Clesceri L.S.** et al. Standard Methods for the Examination of Water and Wastewater (20th ed.). American Public Health Association (APHA). Washington, D.C., 1998, 1325 p.
10. **Su Ch.** et al. A Review on Heavy Metal Contamination on the Soil Worldwide: Situation, Impact and Remediation Techniques. // *Environmental Skeptics and Critics*, 2014, v. 3, № 2, p. 24–38.
11. **Tepanosyan G.** et al. Origin Identification and Potential Ecological Risk Assessment of Potentially Toxic Inorganic Elements in the Topsoil of the City of Yerevan, Armenia. // *Journal of Geochemical Exploration*, 2016, v. 167, p. 1–11.
12. **Gevorgyan G.A.** et al. Environmental Risks of Heavy Metal Pollution of the Soils around Kajaran Town. // *Proceedings of the YSU. Chemistry and Biology*, 2015, № 2, p. 50–55.
13. USEPA Method 3050B: Acid Digestion of Sediments, Sludges and Soils. USA, DC, Washington: Environmental Protection Agency, 1996.
14. **Jiang X.** et al. Potential Ecological Risk Assessment and Prediction of Soil Heavy-Metal Pollution around Coal Gangue Dump. // *Natural Hazards and Earth System Sciences*, 2014, v. 14, p. 1599–1610.